

Lopinavir/Ritonavir

low-PIN-a-veer, rit-ON-uh-veer

Brand Name: Kaletra

Drug Class: Protease Inhibitors

Lopinavir/ritonavir, also known as Kaletra, is a type of medicine called a protease inhibitor (PI). PIs act by blocking protease, a protein that HIV needs to make more copies of itself.

HIV/AIDS-Related Uses

Lopinavir/ritonavir in capsule and oral solution form was approved by the FDA on September 15, 2000, for use with other antiretrovirals in the treatment of HIV infection in adults and children 6 months of age or older. Lopinavir/ritonavir in tablet form was approved by the FDA on October 28, 2005. In March 2006, the capsule form of the medicine was phased out in the U.S. in favor of the new tablet. Lopinavir alone quickly breaks down in the body. Ritonavir slows down this process and increases the time that lopinavir stays in the bloodstream.

This medicine does not cure or prevent HIV infection or AIDS and does not reduce the risk of passing the virus to other people.

Dosage Form/Administration

Lopinavir/ritonavir comes in tablet, oral solution, and capsule forms and is taken by mouth. The tablet can be taken with or without food. The oral solution and capsule forms should be taken with food.

Each film-coated lopinavir/ritonavir tablet contains 200 mg of lopinavir and 50 mg of ritonavir. Every 5 ml of lopinavir/ritonavir oral solution contains 400 mg of lopinavir and 100 mg of ritonavir. Each lopinavir/ritonavir capsule contains 133.3 mg of lopinavir and 33.3 mg of ritonavir.

Recommended Daily Dose

The recommended dose of lopinavir/ritonavir in treatment-experienced adults is 2 tablets (400 mg/100 mg) twice daily taken with or without food or 3 capsules (5 ml) twice daily taken with food. The recommended doses of lopinavir/ritonavir in treatment-naïve adults are 2 tablets (400 mg/100 mg) twice daily taken with or without food, 4 tablets (800 mg/200 mg) once daily taken with or without food.

No dosing adjustment in treatment-naïve patients is

necessary when lopinavir/ritonavir tablets are administered as part of a twice-daily regimen with efavirenz, nevirapine, amprenavir, fosamprenavir, or nelfinavir. A dose increase should also be considered when lopinavir/ritonavir in tablet form is used in combination with efavirenz, nevirapine, fosamprenavir without ritonavir, or nelfinavir in certain treatment-experienced patients. Lopinavir/ritonavir should not be used once daily in combination with efavirenz, nevirapine, amprenavir, fosamprenavir, or nelfinavir. When lopinavir/ritonavir in oral solution or capsule form is used twice daily in combination with efavirenz or nevirapine, the lopinavir/ritonavir dose should be increased. Dosing of lopinavir/ritonavir in children is based on weight.

Some patients may benefit from different doses of lopinavir/ritonavir. Patients should always take lopinavir/ritonavir as prescribed by their doctors.

Contraindications

Individuals should tell a doctor about any medical problems before taking this medicine.

Possible Side Effects

Along with its desired effects, this medicine can cause some serious unwanted effects. Individuals should tell a doctor if they have symptoms of any of the following side effects: disease of the pancreas (symptoms include bloating; chills; constipation; darkened urine; fast heartbeat; fever; indigestion; loss of appetite; pains in stomach, side, or back; yellow eyes or skin), diabetes or high blood sugar (symptoms include blurred vision, dry mouth, fruity mouth odor, increased hunger or thirst, increased urination, fainting, nausea, stomachache, sweating, tiredness, trouble breathing, or unusual weight loss). Other serious side effects include body fat changes, high cholesterol, and increased bleeding in hemophilia patients. Individuals should tell a doctor if they have any of these side effects.



Lopinavir/Ritonavir



Monday - Friday, 12:00 p.m. (Noon) - 4:00 p.m. ET

Possible Side Effects (cont.)

Other side effects may not be serious and may lessen or disappear with continued use of the medicine. Less serious side effects of this medicine include diarrhea, headache, nausea, rash, trouble sleeping, vomiting, and weakness. Diarrhea may occur more frequently in patients taking lopinavir/ritonavir once daily. Individuals should tell a doctor if these side effects continue or are bothersome.

Drug and Food Interactions

A doctor should be notified of any other medications being taken, including prescription, nonprescription (over-the-counter), or herbal medications.

Clinical Trials

For information on clinical trials that involve Lopinavir/Ritonavir, visit the ClinicalTrials.gov web site at <http://www.clinicaltrials.gov>. In the Search box, enter: Lopinavir/Ritonavir AND HIV Infections.

Manufacturer Information

Lopinavir/Ritonavir
Abbott Laboratories
One Hundred Abbott Park Rd
Abbott Park, IL 60064-3500
(800) 633-9110

Kaletra
Abbott Laboratories
One Hundred Abbott Park Rd
Abbott Park, IL 60064-3500
(800) 633-9110

For More Information

Contact your doctor or an AIDSinfo Health Information Specialist:

- Via Phone: 1-800-448-0440 Monday - Friday, 12:00 p.m. (Noon) - 5:00 p.m. ET
- Via Live Help: http://aidsinfo.nih.gov/live_help